

Part VI: Summary of the Risk Management Plan

Summary of Risk Management Plan for BENDAMUSTIN ACTAVIS (BENDAMUSTINE)

This is a summary of the risk management plan (RMP) for Bendamustine Actavis (hereinafter referred to as Bendamustine). The RMP details important risks of Bendamustine, how these risks can be minimised, and how more information will be obtained about Bendamustine's risks and uncertainties (missing information).

Bendamustine's summary of product characteristics (SmPC) and its package leaflet give essential information to healthcare professionals and patients on how Bendamustine should be used.

Important new concerns or changes to the current ones will be included in updates of Bendamustine's RMP.

I. The Medicine and What It is used for

Bendamustine is authorised for (see SmPC for full indication):

- First-line treatment of chronic lymphocytic leukaemia (Binet stage B or C) in patients for whom fludabarine combination chemotherapy is not appropriate
- Indolent non-Hodgkin's lymphomas as monotherapy in patients who have progressed during or within 6 months following treatment with rituximab or a rituximab containing regimen
- Front-line treatment of multiple myeloma (Durie-Salmon stage II with progress or stage III) in combination with prednisone for patients older than 65 years who are not eligible for autologous stem cell transplantation and who have clinical neuropathy at time of diagnosis precluding the use of thalidomide or bortezomib containing treatment

It contains bendamustine as the active substance and it is given by intravenous infusion.

II. Risks Associated with the Medicine and Activities to Minimise or Further Characterise the Risks

Important risks of Bendamustine, together with measures to minimise such risks and the proposed studies for learning more about Bendamustine's risks, are outlined below.

Measures to minimise the risks identified for medicinal products can be:

- Specific information, such as warnings, precautions, and advice on correct use, in the package leaflet and SmPC addressed to patients and healthcare professionals;
- Important advice on the medicine's packaging;
- The authorised pack size — the amount of medicine in a pack is chosen so to ensure that the medicine is used correctly;
- The medicine's legal status — the way a medicine is supplied to the patient (e.g. with or without prescription) can help to minimise its risks.

Together, these measures constitute *routine risk minimisation* measures.

In addition to these measures, information about adverse reactions is collected continuously and regularly analysed so that immediate action can be taken as necessary. These measures constitute *routine pharmacovigilance activities*.

II.A List of Important Risks and Missing Information

Important risks of Bendamustine are risks that need special risk management activities to further investigate or minimise the risk, so that the medicinal product can be safely administered.

Important risks can be regarded as identified or potential. Identified risks are concerns for which there is sufficient proof of a link with the use of Bendamustine. Potential risks are concerns for which an association with the use of this medicine is possible based on available data, but this association has not been established yet and needs further evaluation. Missing information refers to information on the safety of the medicinal product that is currently missing and needs to be collected (e.g. on the long-term use of the medicine).

List of important risks and missing information	
Important identified risks	None
Important potential risks	None
Missing information	None

II.B Summary of Important Risks

The safety information in the proposed Product Information is aligned to the reference medicinal product.

II.C Post-Authorisation Development Plan

II.C.1 Studies Which Are Conditions of the Marketing Authorisation

There are no studies which are conditions of the marketing authorisation or specific obligation of Bendamustine.

II.C.2 Other Studies in Post-Authorisation Development Plan

There are no studies required for Bendamustine.