

## **Part VI: Summary of risk management plan for Fesoterodine Medical Valley 4 and 8 mg forðatöflur (fesoterodine)**

This is a summary of the risk management plan (RMP) for Fesoterodine Medical Valley 4 and 8 mg forðatöflur. The RMP details important risks of Fesoterodine Medical Valley 4 and 8 mg forðatöflur and how more information will be obtained about Fesoterodine Medical Valley 4 and 8 mg forðatöflur risks and uncertainties (missing information).

Fesoterodine Medical Valley 4 and 8 mg forðatöflur summary of product characteristics (SmPC) and its package leaflet give essential information to healthcare professionals and patients on how Fesoterodine Medical Valley 4 and 8 mg forðatöflur should be used.

Important new concerns or changes to the current ones will be included in updates of Fesoterodine Medical Valley 4 and 8 mg forðatöflur's RMP.

### **I. The medicine and what it is used for**

Fesoterodine Medical Valley 4 and 8 mg forðatöflur is authorised for adults, for treatment of the symptoms (increased urinary frequency and/or urgency and/or urgency incontinence) that may occur with overactive bladder syndrome (see SmPC for the full indication). It contains fesoterodine as the active substance, and it is given by oral route.

### **II. Risks associated with the medicine and activities to minimise or further characterise the risks**

Important risks of Fesoterodine Medical Valley 4 and 8 mg forðatöflur, together with measures to minimise such risks are outlined below.

Measures to minimise the risks identified for medicinal products can be:

- Specific information, such as warnings, precautions, and advice on correct use, in the package leaflet and SmPC addressed to patients and healthcare professionals;
- Important advice on the medicine's packaging;
- The authorised pack size — the amount of medicine in a pack is chosen so to ensure that the medicine is used correctly;
- The medicine's legal status — the way a medicine is supplied to the patient (e.g. with or without prescription) can help to minimise its risks.

Together, these measures constitute *routine risk minimisation measures*.

In addition to these measures, information about adverse reactions is collected continuously and regularly analysed so that immediate action can be taken as necessary. These measures constitute *routine pharmacovigilance activities*.

## ***II.A List of important risks and missing information***

Important risks of Fesoterodine Medical Valley 4 and 8 mg forðatöflur are risks that need special risk management activities to further investigate or minimise the risks, so that the medicinal product can be safely taken. Important risks can be regarded as identified or potential. Identified risks are concerns for which there is sufficient proof of a link with the use of Fesoterodine Medical Valley 4 and 8 mg forðatöflur. Potential risks are concerns for which an association with the use of this medicine is possible based on available data, but this association has not been established yet and needs further evaluation. Missing information refers to information on the safety of the medicinal product that is currently missing and needs to be collected (e.g. on the long-term use of the medicine);

<b>List of important risks and missing information</b>	
Important identified risks	None
Important potential risks	None
Missing information	None

## ***II.B Summary of important risks***

The safety information in the Product Information is aligned to the reference medicinal product.

## ***II.C Post-authorisation development plan***

### **II.C.1 Studies which are conditions of the marketing authorization**

There are no studies which are conditions of the marketing authorisation or specific obligation of Fesoterodine Medical Valley 4 and 8 mg forðatöflur.

### **II.C.2 Other studies in post-authorisation development plan**

There are no studies required for Fesoterodine Medical Valley 4 and 8 mg forðatöflur.